

09/647069

TRANSMITTAL LETTER TO THE UNITED STATES

ATTORNEY'S DOCKET NUMBER 48937

DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)


INTERNATIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/EP 99/02275	1 April 1999	6 April 1998
		14 April 1998
		12 August 1998

TITLE OF INVENTION: A PROCESS FOR PREPARING GRIGNARD COMPOUNDS

APPLICANT(S) FOR DO/EO/US Laure BOYMOND, Mario ROTTLAENDER, Gerard CAHIEZ, Paul KNOCHER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. /X/ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. // This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. /X/ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. /x/ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. /X/ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
- a. /X/ is transmitted herewith (required only if not transmitted by the International Bureau).
- b. // has been transmitted by the International Bureau.
- c. // is not required, as the application was filed in the United States Receiving Office (RO/USO).
6. /X/ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. /X/ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
- a. /X/ are transmitted herewith (required only if not transmitted by the International Bureau).
- b. // have been transmitted by the International Bureau.
- c. // have not been made; however, the time limit for making such amendments has NOT expired.
- d. // have not been made and will not be made.
8. /X/ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. /X/ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. // A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern other document(s) or information included:
11. /X/ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. /x/ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. /x/ A FIRST preliminary amendment.
- // A SECOND or SUBSEQUENT preliminary amendment.
14. // A substitute specification.
15. // A change of power of attorney and/or address letter.
16. /x/ Other items or information.
- International Search Report
- International Preliminary Examination Report



U.S. Appl. No. (If Known) INTERNATIONAL APPLN. NO.
PCT/EP99/02275

ATTORNEY'S DOCKET NO.
48937

		CALCULATIONS	PTO USE ONLY
17. /X/ The following fees are submitted			
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):			
Search Report has been prepared by the			
EPO or JPO.....	\$840.00	840.00	
International preliminary examination fee paid to USPTO			
(37 CFR 1.482).....	\$750.00		
No international preliminary examination fee paid to			
USPTO (37 CFR 1.482) but international search fee paid			
to USPTO (37 CFR 1.445(a)(2)).....\$700.00			
Neither international preliminary examination fee			
(37 CFR 1.482) nor international search fee			
(37 CFR 1.445(a)(2)) paid to USPTO\$ 970.00			
International preliminary examination fee paid to			
USPTO (37 CFR 1.482) and all claims satisfied pro			
-visions of PCT Article 33(2)-(4)).....\$96.00			
ENTER APPROPRIATE BASIC FEE AMOUNT =		\$ 840.00	
Surcharge of \$130.00 for furnishing the oath or declaration			
later than // 20 // 30 months from the earliest			
claimed priority date (37 CFR 1.492(e)).			
Claims	Number Filed	Number Extra	Rate
Total Claims	8 -20		X\$18.
Indep. Claims	1 -3		X\$78.
Multiple dependent claim(s) (if applicable)		+270.	
TOTAL OF ABOVE CALCULATION		=	840.00
Reduction of 1/2 for filing by small entity, if applicable.			
Verified Small Entity statement must also be filed			
(Note 37 CFR 1.9, 1.27, 1.28).			
SUBTOTAL		=	840.00
Processing fee of \$130. for furnishing the English			
translation later than // 20 // 30 months from the			
earliest claimed priority date (37 CFR 1.492(f)). +			
TOTAL NATIONAL FEE		=	840.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)).			
The assignment must be accompanied by an appropriate cover			
sheet (37 CFR 3.28, 3.31) \$40.00 per property =			
TOTAL FEES ENCLOSED		= \$	880.00
Amount to be			
refunded: \$			
Charged \$			

a./X/ A check in the amount of \$ 880. to cover the above fees is enclosed.

b./ / Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.

c./X/ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 11-0345. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

KEIL & WEINKAUF
1101 Connecticut Ave., N.W.
Washington, D. C. 20036

SIGNATURE

Herbert B. Keil

NAME

Registration No. 18,967

09/647069

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of)
BOYMOND et al.)

534 Rec'd PCT/PTC 26 SEP 2000
BOX PCT

International Application)
PCT/EP 99/02275)

Filed: April 1, 1999)

For: A PROCESS FOR PREPARING GRIGNARD COMPOUNDS

PRELIMINARY AMENDMENT

Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

Prior to examination, kindly amend the above-identified application as follows:

IN THE CLAIMS

Claim 3, line 1, delete "or 2".

Claim 4, line 1, delete "any of claims 1 to 3" and insert --claim 1--.

Claim 5, line 1, delete "any of claims 1 to 4" and insert --claim 1--.

Claim 8, line 1, delete "any of claims 1 to 5" and insert --claim 1--.

REMARKS

The claims were amended in the preliminary examination. The claims have been amended further to eliminate multiple dependency and to put it in better form for U.S. filing. No new matter is included.

Favorable action is solicited.

Respectfully submitted,

KEIL & WEINKAUF



Herbert B. Keil
Reg. No. 18,967

1101 Connecticut Ave., N.W.
Washington, D.C. 20036
(202)659-0100

534 Rec'd PCT/PTO 26 SEP 2000

A process for preparing Grignard compounds

The invention relates to a process for preparing Grignard compounds of the formula I. The invention additionally relates to compounds of the formula I and to polymer-bound compounds of the formula Ia. The invention further relates to the use of the process for preparing substance libraries and to the use of the compounds of the formulae I and Ia in chemical synthesis.

10


Grignard compounds are valuable intermediates in organic synthesis. They are among the most important classes of compounds in synthetic organic chemistry. Reaction thereof with electrophilic substances allows a wide variety of compounds to be prepared. A large number of syntheses in which Grignard compounds are used are disclosed in the literature (see: Handbook of Grignard-Reagents, Eds. G.S. Silverman, P.E. Rakita, Marcel Dekker, Inc., 1996). Grignard compounds have good reactivity with satisfactory chemoselectivity (see Posner G. H. Org. React., Vol. 22, 1975: 253, Lipshutz et al., Org. React., Vol. 41, 1992: 135, Luh T.-Y. Chem. Res., Vol 24, 1991: 257 or Tamao et al., J. Am. Chem. Soc., Vol. 94, 1972: 4374). DE 196 32 643 describes, for example, the synthesis of intermediates for angiotensin-II inhibitors via a Grignard compound. DE 25 41 438 and DE 19 64 405 describe the synthesis of fragrances by Grignard syntheses, for example for cosmetics.

The German application (file number 19709118.0) describes the synthesis of plant active principles by means of a Grignard compound which has thioalkyl ether or halogen substituents.

They are normally prepared as shown in scheme I, by reacting an appropriate alkyl or aryl halide with metallic magnesium or another magnesium source. The methods for this are known to the skilled worker and can be referred to in the Handbook of Grignard Reagents, Eds. G.S. Silverman, P.E. Rakita, Marcel Dekker, Inc., 1996.

Scheme I: Classical preparation of Grignard compounds

40




However, it is common to all these reactions that they are carried out under rather drastic conditions (temperatures $> 0^{\circ}\text{C}$, usually even at temperatures $> +40^{\circ}\text{C}$). These conditions do not, however, permit other functional groups, such as ester or nitrile moieties, which are able to react with a Grignard compound as electrophile to be retained in the molecule, because oligomerization, reduction or other side reactions would occur under these conditions.

10 Bull. Soc. Chim. Fr. 1967, 1520, Angew. Chem., Vol.81, 1969: 293, J. Organomet. Chem., C21 G, 1971: 33, J. Organomet. Chem. Vol. 113, 1976: 107 and J. Organomet. Chem. Vol. 54, 1973: 123 describe the preparation of aryl Grignard compounds by halogen/magnesium exchange.

15 The conditions and reagents used therein do not, however, allow Grignard compounds with functional groups such as esters, nitriles or amides, which react with an electrophile, to be prepared.

20 Thus, for example, J. Organomet. Chem., Vol. 113, 1976: 107 describes the preparation of 2-pyridylmagnesium bromide (XI) starting from phenylmagnesium bromide (VII) and 2-chloropyridine (VIII) as shown in scheme II.

25 Scheme II: Synthesis of 2-pyridylmagnesium bromide



However, the reaction conditions are such that functional groups in the molecule would be immediately converted.

40 The simple preparation of Grignard compounds which contain functional groups which react with electrophiles has therefore not been disclosed to date.

45


3

It is an object of the present invention to provide a process for synthesizing Grignard compounds which contain other functional groups for many further types of chemical synthesis, which groups are able to react with electrophilic reagents.

5

We have found that this object is achieved by a process for preparing compounds of the general formula I

10



which comprises reacting compounds of the general formula II

15



20

with compounds of the formula R^4MgX (III) at temperatures below $0^\circ C$,

where the substituents and variables in the formulae I, II and

25

$Z = 0, 1$

$X =$ halogen such as Cl, Br, I or R^2

30

$X^a =$ Br, I

A, B, D and E

independently of one another CH, CR^2 , N, P or CR^3

35

$F =$ O, S, NR^6 , CR^2 or CR^3 when $z = 0$, or CH, CR^2 , N, P or CR^3 when $z = 1$,

it being possible for two adjacent variables A, B, D, E or F

40

together to form another substituted or unsubstituted aromatic, saturated or partially saturated ring which has 5 to 8 atoms in the ring and which may contain one or more heteroatoms such as O, N, S, P, and not more than three of the variables A, B, D, E or F being a heteroatom.

45

It is preferred for not more than three of the variables A, B, D, E or F in the compounds of the formulae I, Ia and II to be nitrogen. If $z = 0$, it is possible and advantageous for the ring to contain other heteroatoms such as oxygen or sulfur in addition to the nitrogen or in place of the nitrogen, it being possible for a maximum of one sulfur or oxygen atom to be present in the 5-membered ring. Examples which may be mentioned are 5-membered heterocycles with basic structures such as pyrrole, pyrazole, imidazole, triazole, oxazole, isoxazole, isothiazole, triazole, furazan, oxadiazole, thiooxazole, thiophene or furan. Examples of 6-membered heterocycles which may be mentioned are rings with basic structures such as pyridine, pyrimidine, pyrazine, pyridazine or triazine. It is possible and advantageous, both for $z = 0$ and for $z = 1$, for the only heteroatom in the ring to be a phosphorus atom.

$R^1 =$ COOR², CN, CONR³R^{3'}, halogen

$R^2 =$ substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl, R⁵,

$R^3 =$ hydrogen, substituted or unsubstituted, branched or unbranched -OC₁-C₁₀-alkyl, -OC₃-C₁₀-cycloalkyl, -OC₁-C₄-alkylaryl, -OC₁-C₄-alkylhetaryl, R^{3'} or R⁵,

$R^{3'} =$ hydrogen, substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl, R⁵,

$R^4 =$ substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl or halogen such as Cl, Br, I, preferably Br or I,

$R^5 =$ a solid support, preferably a polymeric protective group,

$R^6 =$ substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl, substituted or unsubstituted, branched or unbranched -(C=O)-C₁-C₁₀-alkyl, -(C=O)-C₃-C₁₀-cycloalkyl, -(C=O)-C₁-C₄-alkylaryl, -(C=O)-C₁-C₄-alkylhetaryl or -SO₂-aryl.

R^1 in the compounds of the formulae I and II is COOR², CN, CONR³R^{3'}, halogen such as F, Cl.

R^2 in the compounds of the formulae I and II is substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl or R^5 .

- 5 Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl chains such as, for example, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

Examples of cycloalkyl radicals in the formula which may be mentioned are substituted or unsubstituted, branched or

- 20 unbranched C_3 - C_{10} -cycloalkyl radicals with 3 to 7 carbon atoms in the ring or ring system such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl, 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl, 1-pentylcyclopropyl, 1-methyl-1-butylcyclopropyl, 1,2-dimethylcyclopropyl, 1-methyl-2-ethylcyclopropyl, cyclooctyl, cyclononyl or cyclodecyl. The cycloalkyl radicals may also contain heteroatoms such as S, N and O in the ring.
- 30 Radicals which may be mentioned as C_1 - C_4 -alkylaryl are substituted and unsubstituted, branched-chain or unbranched-chain C_1 - C_4 -alkylphenyl or C_1 - C_4 -alkylnaphthyl, such as methylphenyl, ethylphenyl, propylphenyl, 1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl, 2-methylpropylphenyl, 1,1-dimethylethylphenyl, methylnaphthyl, ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl, butylnaphthyl, 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or 1,1-dimethylethylnaphthyl.
- 40 Alkylhetaryl radicals which may be mentioned are substituted and unsubstituted, branched-chain or unbranched-chain C_1 - C_4 -alkylhetaryl radicals which contain one or more nitrogen, sulfur and/or oxygen atoms in the ring or ring system.
- 45 R^2 can also be a solid support R^5 (see below for definition of the support).

Suitable substituents of the R^2 radicals mentioned are in principle, apart from ketones or aldehydes, all conceivable substituents, for example one or more substituents such as halogen such as fluorine, chlorine or bromine, cyano, nitro,
 5 amino, hydroxyl, alkyl, cycloalkyl, aryl, alkoxy, benzyloxy, phenyl or benzyl.

R^3 in the substituent R^1 is hydrogen, substituted or unsubstituted, branched or unbranched $-OC_1-C_{10}$ -alkyl,
 10 $-OC_3-C_{10}$ -cycloalkyl, $-OC_1-C_4$ -alkylaryl, $-OC_1-C_4$ -alkylhetaryl, R^3 or R^5 .

$-O$ -Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched $-OC_1-C_{10}$ -alkyl chains (= alkylhydroxamic acids linked via the oxygen). The C_1-C_{10} -alkyl chains in these $-O$ -alkyl radicals have the following meanings:
 15 methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl,
 20 n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl,
 25 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

Examples of $-O$ -cycloalkyl radicals in the radical R^3 which may be mentioned are substituted or unsubstituted, branched or
 30 unbranched $-OC_3-C_{10}$ -cycloalkyl radicals with 3 to 7 carbon atoms in the ring or ring system, where the C_3-C_{10} -cycloalkyl radicals have the following meanings: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl, 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl,
 35 1-pentylcyclopropyl, 1-methyl-1-butylcyclopropyl, 1,2-dimethylcyclopropyl, 1-ethyl-2-ethylcyclopropyl, cyclooctyl, cyclononyl or cyclodecyl. The cycloalkyl radicals may also contain heteroatoms such as S, N and O in the ring.

40

Radicals which may be mentioned as $-O-C_1-C_4$ -alkylaryl are substituted and unsubstituted, branched-chain or unbranched-chain $-O-C_1-C_4$ -alkylaryl, where the C_1-C_4 -alkylaryl radicals have the following meanings: C_1-C_4 -alkylphenyl or C_1-C_4 -alkylnaphthyl
 45 radicals, such as methylphenyl, ethylphenyl, propylphenyl, 1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl, 2-methylpropylphenyl, 1,1-dimethylethylphenyl, methylnaphthyl,

ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl, butylnaphthyl, 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or 1,1-dimethylethylnaphthyl.

- 5 -O-Alkylhetaryl radicals which may be mentioned are substituted and unsubstituted, branched-chain or unbranched-chain -O-C₁-C₄-alkylhetaryl radicals which contain one or more nitrogen, sulfur and/or oxygen atoms in the ring or ring system.
- 10 R³ can also be a solid support R⁵ (see below for the definition of the support) or R^{3'}.

All the abovementioned radicals of the substituent R³ are linked via the oxygen and form so-called hydroxamic acids in the case of
15 the CONR³R^{3'} radical mentioned under R¹, otherwise ethers.

Suitable substituents of the said R³ radicals are in principle, apart from ketones or aldehydes, all conceivable substituents, for example one or more substituents such as halogen such as
20 fluorine, chlorine or bromine, cyano, nitro, amino, hydroxyl, alkyl, cycloalkyl, aryl, alkoxy, benzyloxy, phenyl or benzyl.

R^{3'} in the substituent R¹ is hydrogen, substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl,
25 C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl, R^{3'} or R⁵.

Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl chains such as, for example, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl,
30 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl,
35 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

40 Examples of cycloalkyl radicals in the formula which may be mentioned are substituted or unsubstituted, branched or unbranched C₃-C₁₀-cycloalkyl radicals with 3 to 7 carbon atoms in the ring or ring system such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl,
45 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl, 1-pentylcyclopropyl, 1-methyl-1-butylcyclopropyl, 1,2-dimethylcyclopropyl,

1-methyl-2-ethylcyclopropyl, cyclooctyl, cyclononyl or cyclodecyl. The cycloalkyl radicals may also contain heteroatoms such as S, N and O in the ring.

- 5 Radicals which may be mentioned as C_1 - C_4 -alkylaryl are substituted and unsubstituted, branched-chain or unbranched-chain C_1 - C_4 -alkylphenyl or C_1 - C_4 -alkylnaphthyl, such as methylphenyl, ethylphenyl, propylphenyl, 1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl, 2-methylpropylphenyl,
 - 10 1,1-dimethylethylphenyl, methylnaphthyl, ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl, butylnaphthyl, 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or 1,1-dimethylethylnaphthyl.
 - 15 Alkylhetaryl radicals which may be mentioned are substituted and unsubstituted, branched-chain or unbranched-chain C_1 - C_4 -alkylhetaryl radicals which contain one or more nitrogen, sulfur and/or oxygen atoms in the ring or ring system.
 - 20 $R^{3'}$ can also be a solid support R^5 (see below for definition of the support).
- Suitable substituents of the $R^{3'}$ radicals mentioned are in principle, apart from ketones or aldehydes, all conceivable
- 25 substituents, for example one or more substituents such as halogen such as fluorine, chlorine or bromine, cyano, nitro, amino, hydroxyl, alkyl, cycloalkyl, aryl, alkoxy, benzyloxy, phenyl or benzyl.
 - 30 R^4 in the formula R^4MgX (III) is substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl or halogen such as chlorine, bromine or iodine, preferably bromine or iodine.
 - 35 Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl chains such as, for example, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl,
 - 40 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl,
 - 45 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

Examples of cycloalkyl radicals in the formula which may be mentioned are substituted or unsubstituted, branched or unbranched C₃-C₁₀-cycloalkyl radicals with 3 to 7 carbon atoms in the ring or ring system such as cyclopropyl, cyclobutyl,

- 5 cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl, 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl, 1-pentylcyclopropyl, 1-methyl-1-butylcyclopropyl, 1,2-dimethylcyclopropyl, 1-methyl-2-ethylcyclopropyl, cyclooctyl, cyclononyl or
10 cyclodecyl. The cycloalkyl radicals may also contain heteroatoms such as S, N and O in the ring.

Radicals which may be mentioned as C₁-C₄-alkylaryl are substituted and unsubstituted, branched-chain or unbranched-chain

- 15 C₁-C₄-alkylphenyl or C₁-C₄-alkylnaphthyl, such as methylphenyl, ethylphenyl, propylphenyl, 1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl, 2-methylpropylphenyl, 1,1-dimethylethylphenyl, methylnaphthyl, ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl, butylnaphthyl,
20 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or 1,1-dimethylethylnaphthyl.

Alkylhetaryl radicals which may be mentioned are substituted and unsubstituted, branched-chain or unbranched-chain

- 25 C₁-C₄-alkylhetaryl radicals which contain one or more nitrogen, sulfur and/or oxygen atoms in the ring or ring system.

Suitable substituents of the R⁴ radicals mentioned are in principle, apart from ketones or aldehydes, all conceivable

- 30 substituents, for example one or more substituents such as halogen such as fluorine, chlorine or bromine, cyano, nitro, amino, hydroxyl, alkyl, cycloalkyl, aryl, alkoxy, benzyloxy, phenyl or benzyl.
35 R⁶ is substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₄-alkylaryl, C₁-C₄-alkylhetaryl, substituted or unsubstituted, branched or unbranched -(C=O)-C₁-C₁₀-alkyl, -(C=O)-C₃-C₁₀-cycloalkyl, -(C=O)-C₁-C₄-alkylaryl, -(C=O)-C₁-C₄-alkylhetaryl or
40 -SO₂-aryl.

Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched C₁-C₁₀-alkyl chains such as, for example, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl,

- 45 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl,

10

- 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl,
- 5 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

Examples of cycloalkyl radicals in the formula which may be mentioned are substituted or unsubstituted, branched or

- 10 unbranched C₃-C₁₀-cycloalkyl radicals with 3 to 7 carbon atoms in the ring or ring system such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl, 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl, 1-pentylcyclopropyl, 1-methyl-1-
- 15 butylcyclopropyl, 1,2-dimethylcyclopropyl, 1-methyl-2-ethylcyclopropyl, cyclooctyl, cyclononyl or cyclodecyl. The cycloalkyl radicals may also contain heteroatoms such as S, N and O in the ring.
- 20 Radicals which may be mentioned as C₁-C₄-alkylaryl are substituted and unsubstituted, branched-chain or unbranched-chain C₁-C₄-alkylphenyl or C₁-C₄-alkylnaphthyl, such as methylphenyl, ethylphenyl, propylphenyl, 1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl, 2-methylpropylphenyl,
- 25 1,1-dimethylethylphenyl, methylnaphthyl, ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl, butylnaphthyl, 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or 1,1-dimethylethylnaphthyl.
- 30 Alkylhetaryl radicals which may be mentioned are substituted and unsubstituted, branched-chain or unbranched-chain C₁-C₄-alkylhetaryl radicals which contain one or more nitrogen, sulfur and/or oxygen atoms in the ring or ring system.
- 35 -(C=O)-Alkyl radicals which may be mentioned are substituted or unsubstituted, branched or unbranched -(C=O)-C₁-C₁₀-alkyl chains (= linked via the carbon to which the oxygen is attached by the double bond). The C₁-C₁₀-alkyl chains in these -(C=O)-alkyl radicals have the following meanings: methyl, ethyl, n-propyl,
- 40 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl,
- 45 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl,

1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl,
1-ethyl-2-methylpropyl, n-heptyl, n-octyl, n-nonyl or n-decyl.

Examples of $-(C=O)$ -cycloalkyl radicals in the R^6 radical which may
5 be mentioned are substituted or unsubstituted, branched or
unbranched $-(C=O)-C_3-C_{10}$ -cycloalkyl radicals with 3 to 7 carbon
atoms in the ring or ring system, where the C_3-C_{10} -cycloalkyl
radicals have the following meanings: cyclopropyl, cyclobutyl,
cyclopentyl, cyclohexyl, cycloheptyl, 1-methylcyclopropyl,
10 1-ethylcyclopropyl, 1-propylcyclopropyl, 1-butylcyclopropyl,
1-pentylcyclopropyl, 1-methyl-1-butylcyclopropyl,
1,2-dimethylcyclopropyl, 1-methyl-2-
ethylcyclopropyl, cyclooctyl, cyclononyl or cyclodecyl. The
cycloalkyl radicals may also contain heteroatoms such as S, N and
15 O in the ring.

Radicals which may be mentioned as $-(C=O)-C_1-C_4$ -alkylaryl are
substituted and unsubstituted, branched-chain or unbranched-chain
 $-(C=O)-C_1-C_4$ -alkylaryl, where the C_1-C_4 -alkylaryl radicals have
20 the following meanings: C_1-C_4 -alkylphenyl or C_1-C_4 -alkylnaphthyl
radicals, such as methylphenyl, ethylphenyl, propylphenyl,
1-methylethylphenyl, butylphenyl, 1-methylpropylphenyl,
2-methylpropylphenyl, 1,1-dimethylethylphenyl, methylnaphthyl,
ethylnaphthyl, propylnaphthyl, 1-methylethylnaphthyl,
25 butylnaphthyl, 1-methylpropylnaphthyl, 2-methylpropylnaphthyl or
1,1-dimethylethylnaphthyl.

$-(C=O)$ -Alkylhetaryl radicals which may be mentioned are
substituted and unsubstituted, branched-chain or unbranched-chain
30 $-(C=O)-C_1-C_4$ -alkylhetaryl radicals which contain one or more
nitrogen, sulfur and/or oxygen atoms in the ring or ring system.

All said $-(C=O)$ radicals are linked via the carbon attached to
the oxygen via a double bond.

35 R^6 radicals which may also be mentioned are $-SO_2$ -aryl radicals
such as $-SO_2$ -phenyl or $-SO_2$ -naphthyl. Linkage takes place via the
 SO_2 radical.

40 Suitable substituents for the said R^6 radicals are in principle,
apart from ketones or aldehydes, all conceivable substituents,
for example one or more substituents such as halogen such as
fluorine, chlorine or bromine, cyano, nitro, amino, hydroxyl,
alkyl, cycloalkyl, aryl, alkoxy, benzyloxy, phenyl or benzyl.

45

12

The compounds of the formulae I, Ia and II are aromatic compounds.

The reaction in the process according to the invention is
5 advantageously carried out by reacting the compound II advantageously in an inert, aprotic solvent, for example ethers such as tetrahydrofuran (= THF), diethyl ether, dioxane, dimethoxyethane or methyl tert-butyl ether (= MTB), at temperatures below 0°C, preferably below -10°C, particularly
10 preferably below -15°C, very particularly preferably at -40°C or below, with a compound of the general formula R^4MgX (III) to give a compound of the formula I. It is possible in principle to use all compounds of the formula R^4MgX known to the skilled worker for preparing the Grignard compound, and diisopropylmagnesium or
15 dicyclopentylmagnesium is preferably used. The reaction can generally be carried out in a range from -70°C to 0°C.

Reactions at higher temperatures, for example at 25°C, lead to by-products and thus distinctly lower yields.

20 Under these mild conditions, the halogen/magnesium exchange takes place without the Grignard compounds of the formula II (see above) which are formed reacting with the other functional groups present in the molecule. The compounds react with the
25 electrophile (see examples in Table I) only in the required manner. Conversion with the electrophile greater than 70%, preferably greater than 80%, particularly preferably greater than 85%, very particularly preferably greater than 90%, are possible in this way.

30 The reaction is usually complete within 10 hours, preferably within 5 hours, particularly preferably within 4 hours.

A variant of the process comprises employing not the
35 dialkylmagnesium compound (III with $R^4 = R^2 = X$) directly, but an easily obtainable Grignard compound $X = Hal$ which then, under the reaction conditions, forms in accordance with the Schlenk equilibrium the dialkylmagnesium compound which then undergoes the actual reaction.

40 A particular advantage of this process is that esters of bound alcohols, preferably alcohols bound to polymers, also undergo the halogen/magnesium exchange in the required manner. (Examples in Table II)

45

13

- The linkage of the compounds of the formula I can in this case take place via a solid support (= R⁵) like those known from solid-phase peptide synthesis. Supports which can be used can consist of a large number of materials as long as they are
- 5 compatible with the synthetic chemistry used, it being possible for the size, size distribution and shape of the supports to vary widely depending on the material. Spherical particles are preferred and advantageously have a homogeneous size distribution.
- 10 Examples of suitable solid supports are ceramic, glass, latex, functionalized crosslinked polystyrenes, polyacrylamides, silica gels or resins.
- 15 In order to make attachment of the reactant and elimination of the synthetic product after the synthesis possible, the support must be suitably functionalized or provided with a linker which has an appropriate functional group which makes it possible to attach the compounds according to the invention. Examples of
- 20 suitable and preferred supports and support-linker conjugates are chlorobenzyl-resin (Merrifield resin), Rink resin (Novabiochem), Sieber resin (Novabiochem), Wang resin (Bachem), Tentagel resins (Rapp-Polymere), Pega resin (Polymer Laboratories) or polyacrylamides. Hydroxybenzyl-resin (Wang resin) is particularly
- 25 preferred as support. Very particularly preferred polymeric supports and protective groups are, for example, triphenylmethyl, p-benzyloxybenzyl alcohol, 4-(2',4'-dimethoxyphenyl(hydroxy)methyl)phenoxypolystyrene or 4-(2',4'-dimethoxyphenylmethyl)phenoxypolystyrene.
- 30 The attachment of the compound to the support or polymeric support takes place by reactions which are known to the skilled worker and which are to be found, for example, in the Review by Balkenhohl et al. (Angew. Chem., Vol. 108, 1996: 2436) and the literature cited therein. In the case of Wang resin, the
- 35 attachment can take place, for example, via an ester. The latter can be eliminated from the resin with, for example, trifluoroacetic acid after the synthesis is complete.
- It is possible in this way to utilize the advantages of
- 40 solid-phase synthesis, namely the automatic carrying out and workup of the reaction by simple washing and filtering. It is thus possible to produce substance libraries easily by use of the process according to the invention.
- 45 This means that this reaction is very suitable for generating substance libraries according to the principles of combinatorial chemistry or HSA (Angew. Chem., Vol. 108, 1996: 2436) by first

carrying out the halogen/magnesium exchange on a polymer-linked precursor and then reacting the latter with a large number of electrophiles (in one vessel to generate mixtures).

- 5 After washing and filtering, the target products are then detached from the polymer under conditions suitable for cleavage of the linker bonding.

The compounds according to the invention of the formulae I or Ia
10 (= R⁵ = solid support, preferably polymeric support) can advantageously be used in chemical synthesis as starting materials or intermediates which can be employed in various subsequent reactions. Examples which may be mentioned here are carotenoid, vitamin or active ingredient syntheses, such as
15 active ingredients in the drugs or crop protection sector.

The following examples are intended to illustrate the process without meaning to restrict the method:

20 Examples:

A Preparation of ethyl 4-(α -hydroxybenzyl)benzoate.



























- A solution of 552 mg (2 mmol) of ethyl 4-iodobenzoate in 20 ml of
25 THF was cooled to -40°C and 1.06 mmol of diisopropylmagnesium in methyl tert-butyl ether were added. After 1 h at -40°C 233 mg (2.2 mmol) of benzaldehyde were added. After 3 h, the reaction mixture was hydrolyzed and the org. phase was concentrated. Chromatography of the crude product with 4/1 pentane/ether
30 afforded 460 mg (90%) of the alcohol.

Table I shows the results of analogous reactions with various electrophiles.

- 35 The starting compounds (Grignard compounds) were prepared via iodine/magnesium exchange within half an hour to one hour. The temperature of the reaction solution was between -25 and -40°C. Good conversions were achievable at this temperature. The yields can be increased by using cPent2Mg for the iodine/magnesium
40 exchange reaction (see data in parentheses).

The conversions stated in Table I relate to chemically pure final product. The allylation reactions were carried out in the presence of CuCN x 2 LiCl (10 mol%) (see, for example, Nos. 4-7
45 and 9, 10 and 13).

Table I: Preparation of Grignard compounds and reaction with electrophiles.

	No.	Aryl halide	Electrophile	Product	Yield (%)
5	1		PhCHO		91 (94)
10	2		PhCHO		89
15	3		PhCHO		90
20	4				81
25	5				75
30	6				80 (87)
35	7				79
	8		PhCHO		93
40	9				79
45	10				81

16

No.	Aryl halide	Electrophile	Product	Yield (%)
5 11		PhCHO		94
10 12		HexCHO		74
13				89

15 B. Preparation of Grignard compounds on a polymeric support and reaction with electrophiles.

20 100 mg of Wang resin were mixed with 70 mmol of 4-iodobenzoic acid and 2 ml of THF and cooled to -35°C . 0.7 ml (0.51 mmol) of a 0.73 M solution of diisopropylmagnesium in THF was added dropwise and, after 15 min., 0.7 ml of a 1 M solution of $\text{CuCN}\cdot 2\text{LiCl}$ in THF. Then 0.3 ml of allyl bromide was added and the mixture was stirred for 1 h.

25 Filtration and washing provided the polymer-bound product, which was detached from the polymer under standard conditions (trifluoroacetic acid).

30 The substances listed in Table II were prepared analogously. The yield of free product was usually 90% or more (see data in the table, column 5).

35








40

45

Table II: Preparation of Grignard compounds and reaction with electrophiles on a solid support.

No.	Aryl halide	Electrophile	Product (polymer-bound)	Product (eliminated from the polymer)
1				 >90
2				 >90
3		PhCHO		 >90
4		TosCN		 >90

No.	Aryl halide	Electro- phile	Product (polymer-bound)	Product (eliminated from the poly- mer)
5		TosCN		
6				
7		PhSSPh		
8		PhSSPh		
9		TosCN		

No.	Aryl halide	Electrophile	Product (polymer-bound)	Product (eliminated from the polymer)
10				
11		TosCN		

09/647069

21


534 Rec'd PCT/PTO 26 SEP 2000

We claim:

- 5 1. A process for preparing compounds of the general formula I



which comprises reacting compounds of the general formula II



- 20 with compounds of the formula R^4MgX (III) at temperatures below $0^\circ C$,

where the substituents and variables in the formulae I, II and III have the following meanings:

- 25 $Z = 0, 1$

$X = \text{halogen, } R^2$

- 30 $X^a = \text{Br, I}$

A, B, D and E
independently of one another CH, CR^2 , N, P or CR^3


- 35 $F = \text{O, S, } NR^6, CR^2 \text{ or } CR^3 \text{ when } z = 0, \text{ or CH, } CR^2, \text{ N, P or } CR^3 \text{ when } z = 1,$

- 40 it being possible for two adjacent variables A, B, D, E or F together to form another substituted or unsubstituted aromatic, saturated or partially saturated ring which has 5 to 8 atoms in the ring and which may contain one or more heteroatoms such as O, N, S, P, and not more than three of the variables A, B, D, E or F being a heteroatom,

- 45 $R^1 = COOR^2, CN, CONR^3R^{3'}, \text{ halogen}$

22

- R^2 = substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl, R^5 ,
- 5 R^3 = hydrogen, substituted or unsubstituted, branched or unbranched $-OC_1$ - C_{10} -alkyl, $-OC_3$ - C_{10} -cycloalkyl, $-OC_1$ - C_4 -alkylaryl, $-OC_1$ - C_4 -alkylhetaryl, $R^{3'}$ or R^5 ,
- 10 $R^{3'}$ = hydrogen, substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl, R^5 ,
- 15 R^4 = substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl or halogen,
- R^5 = a solid support,
- 20 R^6 = substituted or unsubstituted, branched or unbranched C_1 - C_{10} -alkyl, C_3 - C_{10} -cycloalkyl, C_1 - C_4 -alkylaryl, C_1 - C_4 -alkylhetaryl, substituted or unsubstituted, branched or unbranched $-(C=O)-C_1$ - C_{10} -alkyl, $-(C=O)-C_3$ - C_{10} -cycloalkyl, $-(C=O)-C_1$ - C_4 -alkylaryl,
- 25 $-(C=O)-C_1$ - C_4 -alkylhetaryl or $-SO_2$ -aryl.
2. A process as claimed in claim 1, which is carried out in an inert aprotic solvent.
- 30 3. A process as claimed in claim 1 or 2, which is carried out at temperatures below -15°C .
4. A process as claimed in any of claims 1 to 3, wherein the reaction to give compounds of the formula I as set forth in
- 35 claim 1 is complete within 10 hours.
5. A process as claimed in any of claims 1 to 4, which is carried out on a solid support ($= R^5$).
- 40 6. A compound of the formula I




45

23

in which R^1 is COOR^2 , CN or $\text{CONR}^3\text{R}^{3'}$ and the other variables and substituents have the meanings stated in claim 1.

7. A compound of the formula Ia



in which variables and substituents have the meanings stated in claim 1, and at least one of the substituents R^2 , R^3 or $R^{3'}$ is a polymeric protective group ($= R^5$).

8. The use of a process as claimed in any of claims 1 to 5 for preparing substance libraries.

A process for preparing Grignard compounds

Abstract

5

The invention relates to a process for preparing Grignard compounds of the formula I. The invention additionally relates to compounds of the formula I and to polymer-bound compounds of the formula Ia. The invention further relates to the use of the

10 process for preparing substance libraries and to the use of the compounds of the formulae I and Ia in chemical synthesis.

15

20

25

30

35

40

45

Declaration, Power of Attorney

Page 1 of 3

0050/048937

We (I), the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

A process for preparing Grignard compounds

the specification of which

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☒ was filed as PCT international application

Number PCT/EP 99/02275

on 01/04/1999

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)–(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
19815078.4	Germany	06 April 1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
19816414.9	Germany	14 April 1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
19836408.3	Germany	12 August 1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Codes, § 119(e) of any United States provisional application(s) listed below.

(Application Number)

(Filing Date)

(Application Number)

(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

2 And we (I) hereby appoint **Messrs. HERBERT. B. KEIL**, Registration Number 18,967; and **RUSSEL E. WEINKAUF**, Registration Number 18,495; the address of both being Messrs. Keil & Weinkauf, 1101 Connecticut Ave., N.W., Washington, D.C. 20036 (telephone 202-659-0100), our attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to sign the drawings, to receive the patent, and to transact all business in the Patent Office connected therewith.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1 - 00 Laure Boymond
NAME OF INVENTOR

16, rue des Celestins
78000 Versailles FR X
France
Citizen of: France
Post Office Address: same as residence

Signature of Inventor

Date

May 04, 1999

2 - 00 Mario Rottländer
NAME OF INVENTOR

Friedrich-Ebert-Str. 117/405
35039 Marburg DE X
Germany
Citizen of: Germany
Post Office Address: same as residence

Signature of Inventor

Date

April 15, 1999

3 - 00 Gerard Cahiez
NAME OF INVENTOR

58, rue de Turbigo
75003 Paris FR X
France
Citizen of: France
Post Office Address: same as residence

Signature of Inventor

Date

May 04, 1999

4 - 00 Paul Knochel
NAME OF INVENTOR

Friedrichsstr. 8
35037 Marburg DE X
Germany
Citizen of: Germany
Post Office Address: same as residence

Signature of Inventor

Date

April 15, 1999